

WZ COUPLINGS OF D-BRANES AND O-PLANES

B. Stefański, jr.

*Department of Applied Mathematics and Theoretical Physics
Cambridge University, Silver Street, Cambridge CB3 9EW, UK
B.Stefanski@damtp.cam.ac.uk*

Abstract In this short note we review the construction and role of Wess-Zumino couplings of Dirichlet branes and Orientifold planes, and show how these combine to give the Green-Schwarz anomaly cancelling terms.

The low energy effective theory of a D-brane has two types of couplings, the Dirac-Born-Infeld (DBI) and Wess-Zumino (WZ) terms. The former describes the coupling of the vector potential and scalars coupling to the NS $\otimes$ NS fields, while the latter gives the coupling of the vector potential and the pullback of the curvature fields to the R $\otimes$ R potentials. The first D-brane WZ term [1] simply states that a D p -brane is charged under $C^{(p+1)}$, the $p + 1$ -form R $\otimes$ R potential

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C^{(p+1)}, \quad (1.1)$$

where μ_p is the charge density of a D-brane and $\mathcal{B}_p$ its worldvolume. It was then shown that D-branes couple to lower dimensional R $\otimes$ R potentials [2] and that a ‘brane within brane’ picture emerged

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C \wedge \text{tr} \exp(i\mathcal{F}/2\pi). \quad (1.2)$$

In the above C represents a formal sum of R $\otimes$ R potentials, $\mathcal{F} = F - B$, F is the worldvolume gauge field and B the NS $\otimes$ NS two form.¹

An R^4 term found in [3] required, as a consequence of duality, the presence of a gravitational WZ coupling [4]. Later the entire WZ action

¹ $\mathcal{F}$, rather than F , is present in the action as the latter quantity is not gauge invariant.

was found using the anomaly inflow mechanism [5, 6] and reads

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C \wedge \text{tr} \exp(i\mathcal{F}/2\pi) \wedge \sqrt{\frac{\hat{A}(R_T)}{\hat{A}(R_N)}}, \quad (1.3)$$

where $\hat{A}$ is the Dirac or A-roof genus and R_T , R_N are the pull-backs of the tangent and normal bundle curvatures to the D-brane world-volume, respectively.

This coupling has a natural interpretation within K-theory [7]. In K-theory there is a natural bilinear pairing of bundles given by the index of the Dirac operator on the tensor product of the two bundles. For E a bundle over a manifold X , with TX the tangent space of X , the map

$$E \rightarrow \text{ch}(E) \sqrt{\hat{A}(TX)}, \quad (1.4)$$

is an isometry with respect to this pairing and the DeRham pairing in $H^*(X)$, the cohomology of X . In fact, following [13], it was shown [14] that D-brane charges are indeed classified by K-theory.

The D-brane WZ couplings were confirmed by string amplitude calculations. In [9] the four-form couplings were computed at tree level. A one-loop amplitude in [10] and tree-level amplitude in [11] confirmed the presence of all the couplings, while in [12] extra, non-anomalous couplings, as well as the normal bundle contributions were determined.

In [8], it was first observed that O-planes too carry gravitational WZ couplings. For consider Type I theory, in which 16 D9-branes, their images and an O9-plane, fill the spacetime. The WZ coupling of the theory is the Green-Schwarz (GS) coupling [15]. This differs from the WZ couplings of 32 D9-branes, indicating that O9-planes too have WZ couplings. By studying one-loop [10], and tree-level [11] scattering amplitudes it was found that the WZ coupling of Op-planes is

$$S_{WZ}^O = -2^{p-4} \mu_p \int_{\mathcal{B}_p} C \wedge \sqrt{\frac{L(R_T/4)}{L(R_N/4)}}, \quad (1.5)$$

where L is the Hirzebruch polynomial. To see that the O9-plane and D9-brane WZ couplings match the GS term consider the following. The massless chiral fields of Type I string theory are a neutral gravitino, a neutral fermion of opposite chirality and $\text{SO}(32)$ fermions. The total anomaly for this theory follows by descent from²

$$I = \frac{1}{2} 2\pi \left(\hat{A}(R) (\text{tr} e^{iR/2\pi} - 2) + \hat{A}(R) \text{tr} e^{iF/2\pi} \right) \Big|_{12\text{-form}}$$

²Note that all of these are real fields.

$$= \pi \frac{1}{(4\pi)^2} (\text{tr} R^2 - \text{tr} F^2) \wedge X_8, \quad (1.6)$$

where

$$X_8 = \frac{1}{(4\pi)^4} \left(\frac{2}{3} \text{tr} F^4 + \frac{1}{12} \text{tr} R^4 + \frac{1}{48} (\text{tr} R^2)^2 - \frac{1}{12} \text{tr} R^2 \text{tr} F^2 \right). \quad (1.7)$$

In the units of [16] the action extracted from the string theory amplitudes [10, 11] is

$$\begin{aligned} S = & -\frac{1}{4\kappa_{10}^2} \int dC^{(2)} \wedge * dC^{(2)} \\ & + \mu_9 \int \left(\frac{2}{(4\pi)^2} C^{(6)} \wedge (\text{tr} R^2 - \text{tr} F^2) + C^{(2)} \wedge X_8 \right). \end{aligned} \quad (1.8)$$

Since $H = dC^{(2)} + \dots$ is gauge invariant and $dC^{(6)} = *dC^{(2)}$ the gauge transformation for $C^{(2)}$ is

$$\delta C^{(2)} = 4\mu_9 \kappa_{10}^2 (\omega_{2,Y}^1 - \omega_{2,L}^1), \quad (1.9)$$

hence (1.8) has an anomalous variation which follows by descent from

$$I_{WZ} = 4\mu_9^2 \kappa_{10}^2 (\text{tr} F^2 - \text{tr} R^2) \wedge X_8. \quad (1.10)$$

The Type I charge density satisfies

$$(\mu_9 \kappa_{10})^2 = \frac{\pi}{2}, \quad (1.11)$$

and hence $I_{WZ} = I$ as required.

O-planes cannot couple to gauge fields so their WZ couplings have to be purely gravitational. Hence it is a consistency check for the D9-brane and GS gauge and mixed couplings to agree.

Acknowledgments

The author would like to thank the conference organisers for financial support and a stimulating programme.

References

- [1] J. Polchinski, *Dirichlet-Branes and Ramond-Ramond Charges*, Phys. Rev. Lett. **75** (1995) 4724, hep-th/9510017;
- [2] M. Li, *Boundary States of D-branes and Dy-Strings*, Nucl. Phys. **B460** (1996) 351, hep-th/9510161; C. Schmidhuber, *D-Brane Actions*, Nucl. Phys. **B467** (1996) 146, hep-th/9601003; M.R. Douglas, *Branes within Branes*, hep-th/9512077;

- [3] C. Vafa and E. Witten, *A One Loop Test of String Dualities*, Nucl. Phys. **B447** (1995) 261, hep-th/9505053;
- [4] M. Bershadsky, C. Vafa, V. Sadov, *D-Branes and Topological Field Theories*, Nucl. Phys. **B463** (1996) 420, hep-th/9511222;
- [5] M.B. Green, J.A. Harvey and G. Moore, *I-Brane Inflow and Anomalous Couplings on D-branes*, Class. Quant. Grav. **14** (1997) 47, hep-th/9605033;
- [6] Y.-K.E. Cheung and Z. Yin, *Anomalies, Branes, and Currents*, Nucl. Phys. **B517** (1998) 69, hep-th/9710206;
- [7] R. Minasian and G. Moore, *K-theory and Ramond-Ramond Charge*, JHEP **9711** (1997) 002, hep-th/9710230;
- [8] K. Dasgupta, D.P. Jatkar, S. Mukhi, *Gravitational Couplings and Z_2 Orientifolds*, Nucl. Phys. **B523** (1998) 465, hep-th/9707224;
- [9] B. Craps and F. Roose, *Anomalous D-brane and Orientifold Couplings from the Boundary State*, Phys. Lett. **445B** (1998) 150, hep-th/9808074;
- [10] J.F. Morales, C.A. Scrucca, M. Serone, *Anomalous Couplings for D-branes and O-planes*, Nucl. Phys. **B552** (1999) 291, hep-th/9812071;
- [11] B. Stefański, jr., *Gravitational Couplings of D-branes and O-planes*, Nucl. Phys. **B548** (1999) 275, hep-th/9812088;
- [12] B. Craps and F. Roose, *(Non-)Anomalous D-brane and O-plane couplings: the Normal Bundle*, Phys. Lett. **450B** (1999) 358, hep-th/9812149;
- [13] For a review see A. Sen, *Non-BPS States and Branes in String Theory*, hep-th/9904207;
- [14] E. Witten, *D-Branes And K-Theory*, JHEP **9812** (1998) 025, hep-th/9810188;
- [15] M.B. Green and J.H. Schwarz, *Anomaly Cancellations in Supersymmetric $D=10$ Gauge Theory and Superstring Theory*, Phys. Lett. **149B** (1984) 117;
- [16] J. Polchinski, *String Theory, Volume 2, Superstring Theory and Beyond*, CUP 1998;